

CREB Working Paper No. 01-16

Education, Health Knowledge, and Maternal Health in Pakistan

Shandana Dar and Uzma Afzal



**Centre for Research in Economics and Business
Lahore School of Economics**

Centre for Research in Economics and Business (CREB)

Naved Hamid
Director CREB

CREB Advisory Board

Shahid Amjad Chaudhry
Rector
Lahore School of Economics

Sohail Zafar
Dean
Faculty of
Business Administration

Azam Chaudhry
Dean
Faculty of Economics

Rana Wajid
Director
Centre for Mathematics
and Statistical Sciences

Muneer Ahmed
Director
Centre for Policy and
Environmental Studies



Lahore School of Economics

Intersection Main Boulevard Phase VI, DHA and Burki Road
Lahore 53200, Pakistan
Tel: 042-36561230; 042-36560936
Email: creb@lahoreschool.edu.pk

CREB Working Paper No. 01-16

Education, Health Knowledge, and Maternal Health in Pakistan

Shandana Dar

MPhil Graduate

Uzma Afzal

Assistant Professor and Research Fellow

© 2016 Centre for Research in Economics and Business
Lahore School of Economics
All rights reserved.

First printing January 2016.

The views expressed in this document are those of the authors and do not necessarily reflect the views of the Centre for Research in Economics and Business or the Lahore School of Economics.

Lahore School of Economics
Intersection of Main Boulevard, Phase VI, DHA, and Burki Road
Lahore 53200, Pakistan
Tel.: +92 42 3656 1230
www.creb.org.pk
creb@lahoreschool.edu.pk

Price: PRs100

Preface

The Centre for Research in Economics and Business (CREB) was established in 2007 to conduct policy-oriented research with a rigorous academic perspective on key development issues facing Pakistan. In addition, CREB (i) facilitates and coordinates research by faculty at the Lahore School of Economics, (ii) hosts visiting international scholars undertaking research on Pakistan, and (iii) administers the Lahore School's postgraduate program leading to the MPhil and PhD degrees.

An important goal of CREB is to promote public debate on policy issues through conferences, seminars, and publications. In this connection, CREB organizes the Lahore School's Annual Conference on the Management of the Pakistan Economy, the proceedings of which are published in a special issue of the Lahore Journal of Economics.

The CREB Working Paper Series was initiated in 2008 to bring to a wider audience the research being carried out at the Centre. It is hoped that these papers will promote discussion on the subject and contribute to a better understanding of economic and business processes and development issues in Pakistan. Comments and feedback on these papers are welcome.

Abstract

The use of maternal health services is a complex behavioral phenomenon related to social and cultural structures, household factors, and women's personal characteristics, such as education and health knowledge. Yet the causal mechanism through which women's education in Pakistan affects their health-seeking behavior is poorly understood. Using data from the 2006/07 Pakistan Demographic Health Survey on women aged 15–49 who had given birth at least once in the three years prior to the survey, this study aims to (i) determine which socioeconomic factors affect maternal healthcare use behavior and (ii) identify the pathway through which the impact of women's education is transmitted to their maternal health-seeking behavior.

The empirical analysis employs a two-step instrumental variable probit model and community fixed effects where women's exposure to mass media is used as an instrument for health knowledge. The results of the analysis indicate that predisposing factors such as women's level of education, their child's birth order, spouse's level of education and occupation, and women's level of empowerment are important determinants of maternal health-seeking behavior in Pakistan. Moreover, women's health knowledge independent of their educational attainment plays an important role in their use of maternal healthcare.

Education, Health Knowledge and Maternal Health in Pakistan

1. Introduction

The World Health Organization (WHO) defines maternal health as the “health of women during pregnancy, childbirth and postpartum period.” It adds that, “while motherhood is a positive and fulfilling experience, for far many women it is associated with suffering, ill health and even death.”

Although women play a principal role in shaping society, they remain at alarmingly high risk of maternal morbidity and mortality. Every life lost in pregnancy and childbirth has multiplicative effects, given that it is women who are responsible for bringing up children and managing household affairs (see Tura & Gebremariam, 2008). These spillovers of maternal health begin from the time a woman becomes pregnant. Decisions made during pregnancy, at the time of delivery, and post-delivery have lifelong implications for mother and child alike. Decreasing maternal mortality is, therefore, one of the most important global agendas of health.

WHO (2013) reports that the maternal mortality ratio (MMR) in Pakistan was 260 per 100,000 live births in 2010. Estimates show that, although Pakistan’s MMR has declined by 47 percent since 1990, the country’s performance remains unsatisfactory compared to other countries in the region. Countries with similar or worse economic conditions, such as Bangladesh and Nepal, have achieved greater success in reducing their MMRs relative to Pakistan.

The Government of Pakistan has made extensive efforts to decrease maternal mortality. The country has some of the most comprehensive mother-and-child healthcare programs with nationwide outreach. Despite this progressive spending, the situation remains dismal. While Pakistan has moved toward meeting most of the Millennium Development Goals, it is likely to miss the goal to reduce maternal mortality to 140 deaths per 100,000 births by 2015 by a large margin.

While the use of maternal health services depends on their provision, the major impediments to improving maternal health tend to be demand-side constraints. Pakistan is a patriarchal society with complex family

structures, and while the decision to seek medical help is made at the individual level, household and community characteristics also play an important role. Therefore, the focus should not only be on increasing the supply of health services, but also on removing the economic, social and cultural barriers that women face in seeking healthcare.

There is an extensive body of literature on the socioeconomic determinants of maternal health in Bangladesh, India and Nepal, but a dearth of such studies on Pakistan. Moreover, the existing studies have limited application for Pakistan at large because they employ data from specific regions or provinces of the country. Most recognize the direct effect of some socioeconomic determinants on maternal health-seeking behavior (age, birth order or previous fetal loss). However, determinants such as education can affect health-seeking behavior through multiple channels. Other branches of health literature, such as child health, have studied at length the pathway through which education affects the health-seeking behavior of an individual. Nevertheless, few studies underpin this causal mechanism in the maternal health literature

This paper is an extension of existing studies that identify key socioeconomic determinants of maternal healthcare utilization by women in developing countries such as Pakistan. Its twofold objective is to (i) determine which socioeconomic factors affect the use of maternal healthcare services by women in Pakistan, and (ii) identify the pathway through which the effect of women's education is transmitted to their maternal health-seeking behavior.

The study draws on data from the Pakistan Demographic Health Survey (PDHS) for 2006/07, which covers approximately 92,340 households. Carried out by the National Institute of Population Studies with the explicit goal of providing much-needed reliable information on maternal and neonatal health in Pakistan, the PDHS gives a comprehensive picture of marriage, fertility preferences, use of family planning methods, and maternal healthcare utilization. It is, therefore, well suited to our purposes.

We use the instrumental variable (IV) technique to estimate the determinants of maternal health behavior; the choice of instrument for the endogenous variable is based on the literature. The results identify women's educational attainment and that of their spouses, women's

health knowledge and their children's birth order as important determinants of maternal health-seeking behavior in Pakistan.

The structure of the paper is as follows. Section 2 provides an overview of the literature on maternal health behavior. Section 3 describes the dataset used, and Section 4 outlines the conceptual framework and econometric methodology. Our empirical findings are presented in Section 5, followed by a conclusion and policy recommendations in Section 6.

2. A Review of the Literature

Maternal health and mortality lie at the heart of development concerns for many developing countries, not only because they reflect the condition of a country's health system, but also because maternal health is a barometer of national development. A vast body of theoretical and empirical literature examines the determinants of maternal health-seeking behavior among women in developing countries (see, for example, Elo, 1992; Bhatia & Cleland, 1995; Celik & Hotchkiss, 2000; Gyimah, Takyi, & Addai, 2006; Mumtaz & Salway, 2007; Sepehri, Sarma, Simpson, & Moshiri, 2008; Amin, Shah, & Becker, 2010; Singh, Rai, Alagarajan, & Singh, 2012).

These studies can be generally categorized into those that (i) identify the supply-side factors determining health-seeking behavior, such as infrastructure quality and the availability of healthcare facilities in a region, and (ii) examine demand-side determinants of maternal healthcare use, such as education, age or women's autonomy. Several recent studies have identified socioeconomic factors as largely determining maternal health-seeking behavior in developing countries. Given the need to improve maternal health in Pakistan, however, there is a surprising dearth of empirical studies in this context. The paper aims to fill this gap and is an extension of the second category.

Among others, a mother's age is a well-established determinant of maternal health behavior (see Elo, 1992; Celik & Hotchkiss, 2000; Das Gupta, Mansuri, Sinha, & Vishwanath, 2007; Amin et al., 2010; Singh et al., 2012). The use of antenatal care and assisted delivery is higher among older women, implying that maternal age serves as a proxy for women's accumulated knowledge (Elo, 1992). At the other end of the spectrum, some studies argue that the use of maternal healthcare is higher among

adolescent women who are emotionally less mature and lack reproductive knowledge (see Amin et al., 2010; Singh et al., 2012).

Numerous studies point to the positive effect of women's schooling on their maternal health-seeking behavior (see Elo, 1992; Celik & Hotchkiss, 2000; Amin et al., 2010; Singh et al., 2012). An interesting observation is that education has a direct and indirect effect on outcome variables. While schooling increases women's health knowledge, the cognitive skills acquired at school increase their ability to assess and assimilate information, and increase their stock of health knowledge even after they leave school. Yet, despite the accepted importance of education for maternal health, most studies have failed to identify the underlying mechanism through which women's schooling affects their health-seeking behavior.

Some exceptions are Elo (1992), Celik and Hotchkiss (2000), and LeVine, LeVine, Rowe, and Schnell-Anzola (2004), who argue that education transforms household dynamics and modifies women's beliefs such that educated women are better able to process knowledge. However, while Elo (1992) and Celik and Hotchkiss (2000) discuss the possible means through which women's education affects their health-seeking behavior, they do not empirically test any such mechanism.

The more recent literature on child health underpins the education–health nexus (see Thomas, Strauss, & Henriques, 1991; Glewwe, 1999; Kovsted, Pörtner, & Tarp, 2002; Webb & Block, 2003; Afzal, 2013). The empirical evidence suggests that much of the education effect translates into health-seeking behavior through the health knowledge that women accumulate with the help of the literacy and numeracy skills they acquire at school.

However, Glewwe (1999), Kovsted et al. (2002), Webb and Block (2003), and Aslam and Kingdon (2012) suggest that, while health knowledge enables women to recognize the need for healthcare, the use of healthcare services helps women acquire additional health knowledge. Health knowledge may, therefore, be an endogenous variable per se if correlated with unobservable individual characteristics as well as with the outcome variable of interest. Studies have tackled this endogeneity issue by using the IV technique, where the most widely used instruments for health knowledge are mass media exposure, maternal education and the presence of close relatives.

To control for this endogeneity issue, we use women's exposure to mass media as an IV for their health knowledge, following the methodology of Glewwe (1999) and Aslam and Kingdon (2012). Glewwe uses household ownership of a radio or television and the availability of newspapers to represent mass media exposure, although the former does not necessarily mean that women have access to the content they convey. Aslam and Kingdon take this a step further and consider a woman exposed to mass media only if she reports watching television.

Even so, this approach to capturing mass media exposure indicates neither the frequency with which women watch television nor the type of programs they view. Based on this, we use the criterion "ever heard a family planning program on radio or television" as a measure of the IV that indicates the specific type of programs women might watch.

Children's birth order is also a key factor in determining whether a woman seeks healthcare. One possible explanation for this is that a woman expecting her first child is more likely to be cautious and rely on medical assistance than women at a higher birth order. Those who have had more children tend to attach less importance to pregnancy, particularly if they have not experienced difficulties during previous pregnancies (Elo, 1992; Celik & Hotchkiss, 2000). Similarly, women who have experienced fetal loss have a higher likelihood of being cautious and using maternal healthcare (Midhet, Becker, & Berendes, 1998).

Prior studies recognize that the patriarchal nature of most Asian societies mean that women's position is multidimensional and intra-household decisions are complex. Male involvement in reproductive healthcare utilization is, therefore, increasingly recognized. Women with educated spouses tend to utilize antenatal care, safe delivery and postnatal care more effectively (Shariff & Singh, 2002; Mullany, Becker, & Hindin, 2007). Similarly, women involved in household decision-making have a higher probability of seeking maternal healthcare (see Furuta & Salway, 2006; Mumtaz & Salway, 2007; Hou & Ma, 2011).

Furuta and Salway (2006) establish that female autonomy is not restricted to control over household finances in the Asian context. They find that women's propensity for discussing family planning with their husbands has a significant effect on the uptake of maternal healthcare. The study captures the extent of women's empowerment using data on discussions

between husband and wife concerning the place of delivery of the last child as a proxy.

Several studies recognize a household's use of health services as a function of its permanent and temporary income, also known as wealth and income, respectively (see, for instance, Celik & Hotchkiss, 2000; Shariff & Singh, 2002; McTavish, Moore, Harper, & Lynch, 2010). However, income tends to fluctuate. Wealth is the more widely used measure because it includes assets that help generate income. Amin et al. (2010) and McTavish et al. (2010) construct a composite wealth index comprising all the durable assets and characteristics of a household. They find that wealthier households use maternal healthcare services more frequently than poorer households. Similarly, larger families imply that a household's existing resources are stretched over more family members, decreasing the affordability – and therefore the use – of maternal healthcare for women (Shariff & Singh, 2002; Mumtaz & Salway, 2007; Singh et al., 2012).

Another important determinant of maternal healthcare at the individual level is geography, which includes community-level factors such as physical access to a hospital or pharmacy or the state of road infrastructure (Elo, 1992; Celik & Hotchkiss, 2000; Gyimah et al., 2006; Gage & Calixte, 2006; Sepehri et al., 2008; Amin et al., 2010). After adjusting for individual-level factors, Gage and Calixte (2006) find that poor roads and lack of transportation significantly reduce the likelihood that a woman will receive antenatal care and four or more antenatal care visits. The availability of a health center within 5 km, however, significantly increases mothers' use of medical assistance.

To some extent, the literature does examine maternal health and its determinants in the Pakistani context (see Midhet et al., 1998; Nisar & White, 2003; Mumtaz & Salway, 2007; Das Gupta et al., 2007; Ali, Bhatti, & Kuroiwa, 2008; Agha & Carton, 2011; Hou & Ma, 2011), but most of these studies are based on simplified assumptions and estimation techniques that render the results questionable. Moreover, their scope is limited because they draw on data from specific regions and provinces of Pakistan. While the recent literature on other countries might provide valuable insight into maternal health, local conditions for developing social policies remain significant.

Most studies on maternal health use only antenatal consultation as a dichotomous dependent variable, although WHO (2006) and Sepehri et al. (2008) argue that full antenatal care comprises tetanus toxoid injections and the frequency of visits to a healthcare facility. In addition, unlike prenatal care and safe delivery, the role of postnatal care as a pillar of safe motherhood is relatively untapped in the literature. Agha and Carton (2011) and Hou and Ma (2011) are two recent studies that employ postnatal care provided by a trained service provider as a component of maternal healthcare services in Pakistan.

Given the backdrop discussed above, the paucity of empirically sound studies on maternal health in Pakistan makes this study an important contribution to the literature. We aim to augment the literature on the determinants of maternal healthcare in Pakistan by identifying the path through which women's education translates into maternal health-seeking behavior – an area that has not been studied in detail thus far.

Lack of information on the quality and availability of infrastructure in a particular region makes it hard to understand the causal effect of such characteristics on women's health-seeking behavior. To take into account such endogeneity problems, we employ community fixed effects and the IV technique. Moreover, the analysis relies not only on the use of antenatal care services, but also on frequency of use, type of delivery, and use of postnatal care as dependent variables.

The dataset used in this study is the PDHS for 2006/07: neither these outcome variables nor the PDHS – which provides a wealth of data – have been used in other studies on fertility in Pakistan.

3. Data and Summary Statistics

The PDHS for 2006/07, a nationally representative, cross-sectional dataset, involves two-stage sampling. In the first stage, a total of 1,000 clusters were identified; in the second stage, 105 households were selected from each sampling point. The survey's purpose is to monitor the population and health situation in Pakistan and track its performance in meeting the Millennium Development Goals.

Unlike conventional demographic and health surveys, the PDHS was designed with the primary objective of obtaining data on maternal and neonatal health. It provides a comprehensive picture of marriage, fertility

preferences, the use of family planning methods and maternal healthcare use. The data was collected from rural as well as urban areas across the four provinces, making it appropriate for a cross-country analysis. As we have already noted, in spite of the extensive information available, very few fertility studies on Pakistan have employed the PDHS 2006/07.

While the survey questionnaire is designed to gather wide-ranging information at both the household as well as individual level, our analysis is conducted at the individual level. The total sample comprises 10,023 women. The working sample includes married women aged 15–49 who gave birth at least once in the three years prior to the survey.¹ The PDHS gives a broad range of information on each birth; to ensure this data is used accurately, we focus on the most recent births closest to the time of interview. This particular sample comprises 4,475 women, 65 percent of who are from rural areas and 34 percent from urban areas.

Table 1 shows that only 39 percent of Pakistani women visited a medical facility for antenatal care at least three times during their last pregnancy. Just over half (51 percent) received at least two tetanus toxoid injections; 38 percent delivered their child at a health facility or under a health professional's care, and only 24 percent utilized postnatal care. Moreover, there is a pronounced difference in the proportion of women who utilized maternal health services in rural and urban areas: only 29 percent of women in rural areas had at least three antenatal visits during pregnancy as opposed to 56 percent in urban areas.

Table 1: Descriptive statistics: mean and standard deviation for overall Pakistan data

Variable	Mean	SD	Observ.
Outcome variables			
Antenatal care			
At least three antenatal visits during pregnancy (= 1)	0.39	0.49	4,418
At least two tetanus toxoid injections during pregnancy (= 1)	0.51	0.50	4,406
Safe delivery (= 1)	0.38	0.49	4,460
Postnatal care (= 1)	0.24	0.43	4,475

¹ We restrict the sample to at least one birth in the three years prior to the survey, given that some women might not respond accurately in connection with births before this interval (see Celik & Hotchkiss, 2000).

Variable	Mean	SD	Observ.
Individual characteristics			
Age			
< 25 (= 1)	0.28	0.45	4,475
25–34 (= 1)	0.52	0.50	4,475
35–49 (= 1)	0.20	0.40	4,475
Education			
None or below primary (= 1)	0.71	0.45	4,475
Primary but below middle (= 1)	0.09	0.29	4,475
Middle but below secondary (= 1)	0.07	0.25	4,475
Secondary and above (= 1)	0.13	0.34	4,475
Literacy skills (= 1)	0.34	0.47	4,475
Health knowledge (= 1)	0.48	0.50	4,461
Mass media exposure			
Heard family planning message on radio or TV in the last month (= 1)	0.37	0.48	4,474
Working woman (= 1)	0.27	0.44	4,471
Child's birth order			
1 (= 1)	0.19	0.39	4,475
2 to 3 (= 1)	0.35	0.48	4,475
4 to 6 (= 1)	0.31	0.46	4,475
7+ (= 1)	0.15	0.36	4,475
Previous fetal loss or stillbirth (= 1)	0.22	0.45	4,475
Previous fetal loss or stillbirth * working woman (= 1)	0.08	0.26	4,471
Planned pregnancy (= 1)	0.74	0.44	4,475
Husband's education			
None or below primary (= 1)	0.41	0.49	4,475
Primary but below middle (= 1)	0.11	0.31	4,475
Middle but below secondary (= 1)	0.15	0.36	4,475
Secondary and above (= 1)	0.32	0.47	4,475
Husband has skilled employment (= 1)	0.53	0.50	4,474
Discussed place of delivery with spouse (= 1)	0.44	0.50	4,475
Husband is a blood relation (= 1)	0.52	0.50	4,471
Wife of household head (= 1)	0.51	0.50	4,474
Household characteristics			
Household wealth (index)			
Low	-1.15	0.33	840
Medium-low	-0.45	0.15	839

Variable	Mean	SD	Observ.
Medium	-0.01	0.11	840
Medium-high	0.39	0.14	840
High	1.21	0.66	839
Number of household members	9.96	5.57	4,475
Community-level characteristics			
Place of residence			
Region			
Urban (= 1)	0.35	0.48	4,475
Rural (= 1)	0.65	0.48	4,475
Province			
Punjab (= 1)	0.40	0.49	4,475
Sindh (= 1)	0.29	0.45	4,475
KP (= 1)	0.20	0.40	4,475
Balochistan (= 1)	0.11	0.32	4,475

Note: (= 1) represents dummy variable; the mean is a proportion of this variable.

Source: Authors' calculations based on PDHS 2006/07.

One of the aims of this study is to identify the direct and indirect mechanisms through which education affects the maternal health behavior of women. The descriptive statistics show that more than 70 percent of women all over Pakistan are illiterate and only 13 percent report having received higher education. We use a series of questions asked as part of the PDHS to measure women's awareness of the risks and problems associated with pregnancy. The descriptive statistics show that less than 50 percent of women are aware of such problems. Interestingly, there is no pronounced difference between rural and urban women in terms of health knowledge.

Next, we construct a wealth index comprising household assets and housing conditions using principal component analysis. This allows us to rank individuals on the basis of their "household scores and divide them in different quintiles, each representing 20 percent of the score between 1 (poorest) and 5 (wealthiest)" (Singh et al., 2012).² Separate t-tests are


² The wealth index is constructed using the following variables: material used to construct roof, walls and flooring; type of sanitation facilities and cooking fuel available; availability of electricity; ownership status of house; ownership of consumer durables (cooler, air conditioner, refrigerator, bicycle, motorcycle, scooter, car, truck, telephone, washing machine, water pump, sewing machine, computer, bed, chairs, cabinets, sofa, camera).

conducted to identify any differences in the individual and household characteristics of women residing in rural and urban Pakistan. The results (available on request) indicate that there is no significant difference in the characteristics of rural and urban women.

4. Conceptual Framework and Estimation Strategy

This study draws on the behavioral model proposed by Anderson (1968) and its modifications. As Figure 1 shows, a household's health-seeking behavior is based on "a sequential and conditional function of an individual's predisposition to use health services, their perceived need to use them and their ability to obtain these services" (Amin et al., 2010, p. 11). The actual use of healthcare is triggered by need during the prenatal and postnatal stages of pregnancy. However, in the absence of such data, we explore the extent to which predisposing and enabling factors affect maternal health behavior.

Figure 1: Conceptual framework for determinants of health-seeking behavior in Pakistan



Source: Amin, Shah, and Becker (2010).

To identify the socioeconomic determinants of maternal healthcare behavior among currently married women in Pakistan, we begin with a simple model:

$$MHB = \beta_0 + \alpha I + \varphi H + \gamma C + \varepsilon \quad (1)$$

where *MHB* is women's maternal health-seeking behavior, *I* represents the vector comprising their individual characteristics, *H* denotes the vector of household characteristics, and *C* represents community characteristics.

4.1. Description of Variables

MHB is the variable of interest, reflecting women's healthcare use during pregnancy. Under the WHO's (2006) recommendations, the following components, when combined, constitute complete maternal healthcare (Table 2).

Table 2: List of dependent variables under MHB

Antenatal care	Dummy variable = 1 if woman has had at least three antenatal visits; = 0 otherwise. Dummy variable = 1 if woman has had at least two tetanus toxoid injections; = 0 otherwise.
Safe delivery	Dummy variable = 1 if delivery took place at a health facility or at home by a doctor, nurse, lady health visitor, auxiliary nurse, midwife, or other health professional; = 0 otherwise.
Postnatal care	Dummy variable = 1 if trained postnatal care was obtained within 42 days of delivery; = 0 otherwise.

Antenatal care is not complete unless women are protected from unhygienic practices and other risks via tetanus toxoid injections. The number of visits to a healthcare facility is also an important factor: by frequently visiting health facilities, women come into direct contact with health professionals, with whom they develop a rapport. Together, these components constitute complete antenatal care.

4.2. Endogeneity and Other Specification Issues

Simple OLS estimation can result in biased estimates due to the feedback effect between certain individual characteristics and women's decision to

seek maternal healthcare, resulting in an endogeneity problem. Unobservable community-level characteristics can also result in biased estimates.

4.2.1. *Women's Education and Maternal Healthcare Behavior*

While the positive association between women's education and their health-seeking behavior is largely undisputed, the mechanism through which this relationship works has not been studied in detail. Apart from the theoretical evidence, the problem is largely methodological.

Equation (1) assumes that women's education has a direct effect on their maternal health-seeking behavior. Schooling can influence women's health behavior by improving their ability to process information efficiently through the cognitive skills (literacy or numeracy) they learn at school. Conversely, the acquisition of health knowledge is not restricted to the period women are enrolled in school – the literacy and numeracy skills they acquire during this time can increase women's health knowledge even after leaving school (Thomas et al., 1991; Glewwe, 1999; Aslam & Kingdon, 2012). This study captures both the direct and indirect effect by adding the following pathways of education as control variables:

- Education → literacy skills → maternal health behavior
- Education → literacy skills → health knowledge → maternal health behavior

The literacy skills acquired at school enable women to assess and assimilate information successfully, and increase their health knowledge, which affects their health-seeking behavior (Glewwe, 1999). Maternal health knowledge is itself a potentially endogenous variable. While health knowledge helps identify the need to avail maternal health services, women who seek healthcare are likely to have greater health knowledge. Thus, the causality runs in both directions.

This endogeneity issue can be resolved by using the IV technique. Ideally, the instrument Z_i should contain covariates that have strong explanatory power for the specific endogenous variable, but no correlation with ε , that is, $\text{Cov}(X_i, Z_i) \neq 0$ and $\text{Cov}(\varepsilon_i, Z_i) = 0$.

An important source of information and health knowledge for women is their exposure to mass media (Thomas et al., 1991; Glewwe, 1999; LeVine et al., 2004; Aslam & Kingdon, 2012). This study instruments for health knowledge by using the variable “mass media exposure” as an IV.

The PDHS provides information on households’ ownership of a radio or television, along with the type of health message delivered to women through these media. Based on the availability of such factors, this study considers a woman having been exposed to mass media only if she reports having “heard a family planning message on radio or television last month.”

We assume that this particular measure of mass media increases awareness among women, improves their health knowledge, and affects their health-seeking behavior. The instrument is informative because listening to a family planning message on the radio or television is unlikely to affect women’s health-seeking behavior through any other mechanism but increased health knowledge. Moreover, it is highly improbable that women would listen to family planning messages on the radio or television because they had received maternal healthcare. This makes mass media exposure a valid instrument

4.2.2. Omitted Variable Problem

Short-run fluctuations in a household’s resources, represented by its income or income per capita, are an important determinant of the ability and willingness to pay for healthcare services. Following the pattern of the USAID-sponsored demographic and health surveys, the PDHS 2006/07 does not have any information on household income, but does provide extensive data on ownership of household assets.

Based on the literature and availability of factors in the data, household income is gauged using a wealth index (an index of durable assets owned by the household) as an independent variable. It is important to note that the wealth index does not include ownership of a radio or television. This is because the data shows there is a strong correlation between households that own either a radio or television and women who report having heard a family planning message through either medium. For instance, Table 3 shows that over 90 percent of women from households that own a television reported listening to a family planning message in the last month. Including ownership of these consumer durables would,

therefore, decrease the explanatory power of the endogenous variable since women's mass media exposure would no longer be an informative instrument.

We also use the level of education attained by women's spouses and their occupation as independent variables. These covariates closely reflect spouses' earnings and control for the relative socioeconomic status of a household.

Table 3: Proportion of women from households that own a radio or television

Ownership status of household	Number	Percentage
Owens a television	1,421	90.68
Owens a radio	616	39.39

Source: Authors' calculations based on PDHS 2006/07 data.

Although community-level covariates such as the quality of local health infrastructure cannot be included in the model (due to data unavailability), they are important determinants of healthcare use and can result in biased coefficients unless controlled for. Elo (1992) notes that better-educated women might belong to a particular community that has more schools, for instance. This can confound the relationship between women's education and their health-seeking behavior.

To eliminate the omitted variable bias caused by community characteristics, we estimate a cluster fixed effects model. Fixed effects at a smaller level eliminate all the characteristics common to women in a single cluster, thus avoiding omitted variable bias.

4.3. Estimation Strategy

Given that the predicted probabilities of the outcome variables are bound between 0 and 1, we use a probit model to assess the impact of socioeconomic factors on the uptake of maternal health services. Since health knowledge is potentially endogenous, the model is estimated using the IV technique.

The first stage of two-stage least squares involves ordinary least squares (OLS) to predict the probability of health knowledge. The second stage is the probit model, based on the binary dependent variables.

Stage 1 is represented by the following equation:

$$\text{Health knowledge}_i = \psi_0 + \Sigma\alpha I_1 + \Sigma\phi H_j + \Sigma\gamma C_c + \lambda \text{mass media exposure}_i + \varepsilon_i \quad (2)$$

Stage 2 is represented by the following equation:

$$MHB_i = \beta_0 + \Sigma\alpha \hat{I}_i + \phi \text{health knowledge}_i + \Sigma\phi H_j + \Sigma\gamma C_c + \varepsilon_i \quad (3)$$

Vector I captures individual-level characteristics; the household's environment is captured by vector H . C represents community characteristics, *health knowledge* (measured by women's awareness of the risks and complications of pregnancy) is the endogenous variable, and *mass media exposure* is the IV. This selection of variables is drawn from the choice of variables in the literature and the corresponding data available in the PDHS 2006/07.

In addition, we estimate the following fixed effects logit model, using the IV method:

$$MHB_{ic} = \beta_0 + \Sigma\alpha_i + \Sigma\phi_j + \theta_c + \varepsilon_{ic} \quad (4)$$

Equation (4) represents fixed effects at the cluster level (the primary sampling unit). This model controls for the distance to a health facility and the quality of local health infrastructure as well as other unobservable heterogeneity within sampling units, which cannot be controlled for by the covariates included in the analysis. The variable θ captures the unobserved heterogeneity constant across communities, that is, all unobserved factors (constant among sampling units) that affect maternal health-seeking behavior. It is important to note that the fixed effects model is based on the IV technique.

5. Results and Empirical Findings

The econometric specification we follow uses all the possible variables given in Figure 1 that could affect women's maternal health-seeking behavior in Pakistan.

5.1. First-Stage Results

In the first stage of the two-step probit model, the endogenous covariate is regressed on the IV using simple OLS (see Table A1 in the Appendix). Note that this stage remains the same for all the dependent variables used.

To establish that mass media exposure is an informative instrument, we carry out an Angrist-Prischke F-test. The first-stage F-statistic value is 25.30, which is greater than 10, allowing us to reject the null hypothesis (i.e., that the IV has no explanatory power). Moreover, the significant coefficient of the IV in the first-stage regression indicates that mass media exposure strongly influences women's health knowledge. It is important to note that the programs disseminating information on family planning are public service programs with nationwide outreach. Along with private channels, they are also broadcasted on national channels, irrespective of the type of area or village to which the target audience might belong. This makes mass media exposure an exogenous variable.

The second stage, based on the probit model, uses the predicted endogenous covariate obtained from the first stage as an explanatory variable, along with other individual and household-level characteristics to identify which factors determine women's maternal healthcare use. The estimations are conducted using data for Pakistan overall, controlling for regional and provincial differences among the three provinces of Punjab, Sindh, and Khyber Pakhtunkhwa (KP). Further, to control for community-specific factors such as the quality of local health infrastructure, we employ cluster fixed effects. In all cases, we control for heteroskedasticity of unknown form by implementing robust standard errors at the cluster level.

The explanatory variables are classified as individual characteristics, household characteristics, and community characteristics. Women's individual characteristics, their household characteristics, and community characteristics are regressed separately on four binary dependent variables: (i) made at least three antenatal visits to a healthcare facility (yes = 1, no = 0), (ii) received at least two tetanus toxoid inoculations (yes = 1, no = 0), (iii) had a safe delivery (yes = 1, no = 0), and (iv) received postnatal care (yes = 1, no = 0).

5.2. Second-Stage Results

Table 4 reports the estimated parameters of the data for Pakistan overall for each outcome variable. The model also addresses the endogeneity problem of reverse causality. The probability derivatives or marginal effects of these estimated parameters, which indicate the per unit change in the independent variable and its effect on the outcome probability, are estimated separately and are available on request.

In most cases, the signs of the variables are consistent with the literature on women's maternal health-seeking behavior.

Women's own characteristics include their age, categorized into age cohorts. Column 1 suggests that women aged 35–49 are at least three times as likely to receive antenatal care than women under the age of 25. Similarly, women aged 35–49 have a greater probability of receiving at least two tetanus toxoid inoculations during pregnancy, safe delivery care, and postnatal care compared to women under 25. This effect conforms to Elo (1992), who finds that maternal healthcare use in Peru was higher among older women.

The coefficients for different categories of women's educational attainment show that women with middle and higher education are more likely to visit a healthcare facility seeking antenatal care at least three times during pregnancy compared to uneducated women. Interestingly, women who have received a formal education, but not completed middle school, also have a greater probability of making at least three antenatal visits. These results remain consistent for other aspects of maternal health-seeking behavior, such as tetanus toxoid inoculations, safe delivery, and postnatal care. This implies that all levels of education have a significantly positive effect on women's maternal health-seeking behavior (see Elo, 1992; Amin et al., 2010; Singh et al., 2012).

As mentioned in Section 4, women's education has an indirect effect on their maternal health-seeking behavior. Formal schooling enables women to assess and assimilate information more efficiently, which increases their health knowledge and thus affects the outcome variable. Health knowledge itself is, therefore, an important explanatory variable of health-seeking behavior.

Table 4: Probit results for women's maternal healthcare behavior

Explanatory variable	Antenatal care		Safe delivery	Postnatal care
	(1)	(2)	(3)	(4)
Individual char.				
Age (years)				
< 25 (ref.)				
25–34	0.0904 (0.0595)	-0.0359 (0.0577)	0.0823 (0.0590)	0.0800 (0.0626)
35–49	0.333*** (0.0875)	0.219*** (0.0824)	0.360*** (0.0824)	0.221** (0.0879)
Education				
None or less than primary (ref.)				
Primary but below middle	0.372*** (0.0782)	0.472*** (0.0782)	0.324*** (0.0770)	0.233*** (0.0821)
Middle but below secondary	0.499*** (0.0872)	0.563*** (0.0899)	0.449*** (0.0868)	0.337*** (0.0862)
Secondary and above	0.526*** (0.0885)	0.440*** (0.0877)	0.683*** (0.0866)	0.424*** (0.0851)
Health knowledge	3.984*** (0.528)	4.353*** (0.538)	1.684*** (0.555)	1.987*** (0.580)
Working woman	-0.620*** (0.0940)	-0.668*** (0.0946)	-0.385*** (0.0969)	-0.385*** (0.104)
Child's birth order				
1 (reference)				
2 to 3	-0.432*** (0.0705)	-0.252*** (0.0622)	-0.279*** (0.0668)	-0.234*** (0.0706)
4 to 6	-0.573*** (0.0852)	-0.406*** (0.0808)	-0.504*** (0.0828)	-0.371*** (0.0853)
7+	-0.896*** (0.117)	-0.709*** (0.106)	-0.583*** (0.118)	-0.558*** (0.127)
Previous fetal loss or stillbirth	-0.363*** (0.0872)	-0.439*** (0.0827)	-0.137 (0.0857)	-0.162* (0.0852)
Previous fetal loss or stillbirth * working woman	0.384*** (0.115)	0.206* (0.118)	0.233** (0.116)	0.122 (0.129)
Wanted pregnancy	0.0727 (0.0662)	0.318*** (0.0640)	0.126* (0.0668)	0.215*** (0.0716)
Husband's education				
None or less than primary (ref.)				
Primary but below middle	-0.0485 (0.0766)	-0.0122 (0.0735)	0.120 (0.0757)	0.0604 (0.0816)
	0.150**	0.139**	0.123*	0.0961

Explanatory variable	Antenatal care		Safe delivery	Postnatal care
	(1)	(2)	(3)	(4)
Middle but below secondary	(0.0696)	(0.0668)	(0.0672)	(0.0761)
Secondary and above	0.249*** (0.0609)	0.200*** (0.0593)	0.253*** (0.0598)	0.279*** (0.0619)
Husband has skilled employment	0.226*** (0.0502)	0.127** (0.0499)	0.0169 (0.0516)	0.0983* (0.0563)
Discussed place of delivery with spouse	0.157** (0.0681)	-0.0729 (0.0701)	0.342*** (0.0700)	0.353*** (0.0745)
Husband is a blood relation	-0.178*** (0.0479)	-0.0659 (0.0461)	-0.0517 (0.0476)	-0.00522 (0.0518)
Wife of household head	-0.117** (0.0550)	-0.183*** (0.0504)	-0.147*** (0.0553)	-0.159*** (0.0603)
Household char.				
Household wealth (index)	0.139*** (0.0309)	0.0461 (0.0319)	0.179*** (0.0316)	0.140*** (0.0329)
Number of household members	-0.00941* (0.00524)	-0.00703 (0.00503)	-0.00950* (0.00547)	-0.00682 (0.00521)
Community char.				
Place of residence				
Region				
Urban (ref.)				
Rural	-0.245*** (0.0565)	-0.0929 (0.0575)	-0.312*** (0.0611)	-0.268*** (0.0617)
Province				
Punjab (ref.)				
Sindh	-0.756*** (0.149)	-1.244*** (0.153)	-0.0701 (0.161)	-0.110 (0.165)
KP	-0.0514 (0.0668)	-0.0267 (0.0743)	0.172** (0.0790)	-0.102 (0.0823)
Constant	-1.491*** (0.194)	-1.350*** (0.193)	-0.934*** (0.207)	-1.658*** (0.222)
Number of observations	4,133	4,125	4,172	4,186

Notes: * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$. Robust standard errors at cluster level are given in parentheses below estimates.

Col. 1 = min. three antenatal visits to health facility, col. 2 = received at least two tetanus toxoid injections, col. 3 = gave birth at health facility/by trained health service providers, col. 4 = received trained postnatal care.

Source: Authors' calculations.

By controlling for education, this study intends to isolate the impact of women's health knowledge on their health-seeking behavior. The positive and statistically significant coefficient of the health knowledge variable

for all outcome variables supports this hypothesis, implying that women who are more aware of the issues and complications associated with pregnancy have a higher probability of seeking maternal healthcare (see columns 1–4 in Table 4). It is interesting to note that health knowledge has a more pronounced effect on complete antenatal care use than safe delivery and postnatal care.

The literature has established that women's employment status is associated with their empowerment and better healthcare-seeking behavior. Contrary to other studies, our results show that working women have a smaller probability of seeking antenatal care than women who are not working. This could be due to the former's financial status and nature. Work status may be picking up the effect of other socioeconomic characteristics of working women, for example, the combined effect of the household's financial standing, the nature of the woman's job, and the extent to which she is involved in intra-household decision making.

Among the 27 percent of women of the working sample who are part of the labor force, 79.5 percent are illiterate. Such low levels of education translate into lower earnings for these women. Moreover, of all the women who are working, 38 percent belong to the least wealthy households and only 7 percent belong to the wealthiest households (see Table A2). These statistics show that most working women fall within the lower income strata of the sample.

Table A3 reiterates this by showing that, of the 27 percent of women who are working, only 14 percent have a skilled job while 86 percent are low-paid, unskilled workers. The estimations, therefore, pick up the effect of working women with lower earnings who belong to poor households and cannot afford to seek healthcare.

In Table 4, the coefficients in columns 1, 2, 3, and 4 for all birth orders show that women expecting their first child have a greater probability of making at least three visits to seek antenatal care compared to those with a higher birth order. Interestingly, this effect becomes large for all components of maternal healthcare as birth order increases. This is in accord with the literature and suggests that women with more children tend to attach less importance to subsequent pregnancies.

Contrary to other findings on maternal health-seeking behavior, our results indicate that women who have experienced a fetal loss or stillbirth

have less likelihood of seeking antenatal care, tetanus inoculations, and postnatal care compared to women with no maternal history. Given the unexpected sign of the result, we investigate the interaction of the variable with women's working status. The estimated parameters reported in column 3 show that whether a woman has a history of fetal loss or stillbirth does not have any effect on her probability of seeking safe delivery care.

However, when this variable interacts with the working women variable, the coefficient becomes positive and statistically significant for antenatal visits, tetanus inoculations, and safe delivery. The interaction term is constructed to gauge the impact of maternal history along with the household's socioeconomic status on the outcome variables. These results indicate that the likelihood of a woman visiting a health facility to seek antenatal care at least three times during pregnancy, of receiving tetanus inoculations, and of utilizing safe delivery care is higher if she is a working woman and has experienced fetal loss or stillbirth, keeping all other factors constant.

The literature suggests that women going through a planned pregnancy are more cautious and tend to attach greater importance to the child expected. The results for the variable planned pregnancy support the literature: it has a positive and significant effect on tetanus inoculations and use of safe delivery and postnatal care by women. This implies that a woman facing a planned pregnancy is more cautious and, therefore, has a higher probability of receiving maternal healthcare compared to a woman going through an unexpected pregnancy (see Ahmed & Mosley, 2002; Sepehri et al., 2008).

The level of education attained by a woman's husband is hypothesized to reflect the household's economic wellbeing as it translates into higher earnings. At the same time, the husband's education reflects his perceptions and preferences with respect to modern medicine. The coefficient of the spouse's primary education is statistically insignificant for all the outcome variables, indicating that primary education alone is not enough to ensure higher earnings or modified attitudes toward maternal healthcare. However, women whose spouses have completed at least middle or higher schooling are more likely to seek maternal healthcare than women with illiterate spouses. These results are consistent with the literature, which argues that a higher level of spousal

education translates into greater household income and increases women's probability of seeking maternal healthcare.

Another proxy for household economic wellbeing is whether a woman's spouse has skilled employment. The effect of this variable is also consistent with the literature: it shows that the probability of seeking antenatal care and receiving tetanus inoculations and postnatal care is higher for women whose spouses have skilled employment compared to those whose spouses have unskilled jobs. However, our results indicate that skilled employment in this context has no effect on women's likelihood of receiving safe delivery care.

This study uses spousal discussion of the place of delivery as a proxy for women's involvement in the decision-making process. A woman's ability to make important decisions and influence her personal environment emerges as a strong determinant of maternal health behavior in the literature (see Furuta & Salway, 2006; Mumtaz & Salway, 2007; Hou & Ma, 2011). In Pakistan, however, where family structures are usually complex and most women live in a joint family system, a woman's ability to make independent decisions is not an adequate measure of her autonomy. The variable spousal discussion is, therefore, a viable proxy for women's involvement in the decision-making process.

Spousal discussion has a positive and significant effect on the likelihood of making at least three antenatal visits and seeking safe delivery and postnatal care. This implies that women who discuss such decisions with their spouse have a higher probability of seeking maternal healthcare, after controlling for all other individual and household characteristics (see Furuta & Salway, 2006). Conversely, spousal discussion concerning the place of delivery does not influence women's decision to get at least two tetanus inoculations.

We also include the variable "wife of household head" to see if this has any effect on women's health-seeking behavior. Contrary to expectations, a woman who is the household head's wife has less probability of visiting a healthcare facility at least three times during pregnancy. The effect of this variable remains the same for tetanus inoculations, safe delivery, and postnatal care. Consanguineous marriages are very common in Pakistan and 52 percent of women in the working sample are married to their cousins. However, as the results in Table 4 show, controlling for other factors, consanguinity does not influence any outcome variable.

Household wealth and the number of household members are included in the model to capture household wellbeing. A wealth index comprising household conditions and ownership of durable assets is created using principal component analysis. The coefficient shows that, *ceteris paribus*, women from wealthier households have a greater probability of receiving maternal healthcare.

The coefficients for the number of household members in relation to making at least three antenatal visits and safe delivery care are negative, implying that being part of a larger household has a strong adverse impact on women's antenatal care and safe delivery care utilization. This finding is in accord with the literature, which suggests that women from larger households have fewer resources at their disposal and are subject to more congestion and thinly stretched scarce resources compared to women in smaller households.

However, the size of a household does not influence the probability of a woman receiving tetanus inoculations. This is because, unlike other maternal health-seeking practices, tetanus inoculations do not necessarily involve mobility or incur any cost. Women visited by lady health workers are more likely to receive tetanus inoculations than those who are not. Unfortunately, there is not enough data in the PDHS to investigate behavioral differences in maternal health seeking caused by services delivered by lady health workers.

The results of Table 4 show that women living in rural areas are less likely to receive maternal healthcare than their urban counterparts. This is not surprising: urban areas have greater access to healthcare facilities than rural areas. However, there is no significant difference between rural and urban women's probability of receiving at least two tetanus inoculations. An interaction term is created for each variable that is statistically different between rural and urban areas (e.g., planned pregnancy) with the rural dummy variable. The effect of these interaction terms is insignificant for all the dependent variables and is not, therefore, included in the analysis.

At the provincial level, women in Sindh have a smaller probability of receiving complete antenatal care than women in Punjab. Fewer women in KP use safe delivery care compared to women in Punjab, after controlling for all other factors. Interestingly, there is no significant variation between women in Punjab and KP in terms of receiving complete antenatal and postnatal care.

5.3. Community Fixed Effects

The socioeconomic conditions of an area, cultural factors, and community-specific characteristics – such as the distance to the nearest health facility, the availability of public transport, and the quality of local health infrastructure – also affect the variable of interest and outcome variable. However, based on the availability of factors in the PDHS, these characteristics cannot be included in the analysis.

There is growing evidence in the literature that community-specific characteristics, such as the number of schools in a community, may affect the level of education attained by women as well as by their spouses. This is either because better-educated men and women belong to regions with better socioeconomic conditions or because educated people are more likely to migrate to areas with better economic opportunities.

To address this issue of omitted variable bias and isolate the impact of education, we employ community fixed effects at the cluster level in a separate regression analysis. On average, there are about 10 to 15 households in a primary sampling unit. Households within a primary sampling unit are located in the same vicinity so that observable and unobservable characteristics can be taken into account in the fixed effects model.

Table 5 provides fixed effects estimates at the cluster level for women's maternal health-seeking behavior. In the case of at least three antenatal visits made to a health facility, the effect of all levels of women's education remains significant. However, there is a substantial decline in coefficient size, indicating that cluster-specific characteristics affect women's education in the IV probit model. Similarly, the effect of all levels of education remains significant, but when the size of the coefficient declines for safe delivery care and postnatal care utilization, these cluster-specific characteristics are controlled for via fixed effects.

Table 5: Cluster fixed effects for women's maternal healthcare use behavior

Explanatory variable	Antenatal care		Safe delivery	Postnatal care
	(1)	(2)	(3)	(4)
Individual char.				
Age (years)				
< 25 (ref.)				
25–34	0.00897 (0.0279)	-0.0110 (0.0308)	0.00725 (0.0232)	0.0144 (0.0207)
35–49	0.0521 (0.0376)	0.0167 (0.0413)	0.0779** (0.0313)	0.0363 (0.0281)
Education				
None or less than primary (ref.)				
Primary but below middle	0.0920*** (0.0357)	0.0639 (0.0395)	0.0802*** (0.0296)	0.0464* (0.0265)
Middle but below secondary	0.0952** (0.0407)	0.104** (0.0450)	0.144*** (0.0340)	0.0802*** (0.0304)
Secondary and above	0.0794* (0.0417)	0.0800* (0.0453)	0.163*** (0.0341)	0.103*** (0.0304)
Health knowledge	0.812*** (0.257)	0.942*** (0.287)	0.388* (0.212)	0.378** (0.189)
Working woman	-0.0639** (0.0292)	-0.0518 (0.0317)	-0.0663*** (0.0238)	-0.0630*** (0.0213)
Child's birth order				
1 (reference)				
2 to 3	-0.104*** (0.0276)	-0.0700** (0.0304)	-0.0736*** (0.0229)	-0.0602*** (0.0205)
4 to 6	-0.166*** (0.0348)	-0.139*** (0.0387)	-0.156*** (0.0291)	-0.0941*** (0.0260)
7+	-0.218*** (0.0454)	-0.194*** (0.0500)	-0.138*** (0.0378)	-0.0998*** (0.0337)
Previous fetal loss or stillbirth	-0.0379 (0.0313)	-0.0428 (0.0345)	-0.0116 (0.0262)	-0.0105 (0.0235)
Previous fetal loss or stillbirth * working woman	0.0748 (0.0504)	0.00330 (0.0557)	0.0517 (0.0418)	-0.00291 (0.0375)
Wanted pregnancy	-0.0160 (0.0240)	0.0696* (0.0271)	0.0303 (0.0202)	0.0421** (0.0179)

Explanatory variable	Antenatal care		Safe delivery	Postnatal care
	(1)	(2)	(3)	(4)
Husband's education				
None or less than primary (ref.)				
Primary but below middle	-0.0577* (0.0333)	-0.0258 (0.0362)	0.00177 (0.0274)	-0.00683 (0.0246)
Middle but below secondary	0.0263 (0.0311)	0.0208 (0.0350)	0.0214 (0.0261)	0.0104 (0.0233)
Secondary and above	0.0650** (0.0274)	0.0609** (0.0303)	0.0791*** (0.0228)	0.0929*** (0.0204)
Husband has skilled employment	0.0390* (0.0221)	0.00171 (0.0245)	-0.0179 (0.0182)	0.00329 (0.0163)
Discussed place of delivery with spouse	0.0944** (0.0281)	0.0147 (0.0312)	0.121*** (0.0229)	0.108*** (0.0203)
Husband is a blood relation	-0.0116 (0.0189)	-0.0242 (0.0209)	-0.0149 (0.0158)	-0.0197 (0.0141)
Wife of household head	-0.00425 (0.0226)	-0.0125 (0.0248)	-0.0416** (0.0187)	-0.0333** (0.0167)
Household char.				
Household wealth (index)	0.0314** (0.0135)	0.0171 (0.0150)	0.0328*** (0.0113)	0.0.235** (0.00998)
Number of household members	-0.00108 (0.00215)	-0.00357 (0.00238)	-0.00164 (0.00179)	-0.00192 (0.00161)
Number of observations	4,054	4,046	4,094	4,108
Adjusted R-squared	-0.778	-1.080	-0.318	-0.319

Notes: * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$. Robust standard errors at cluster level are given in parentheses below estimates.

Col. 1 = min. three antenatal visits to health facility, col. 2 = received at least two tetanus toxoid injections, col. 3 = gave birth at health facility/by trained health service providers, col. 4 = received trained postnatal care.

Source: Authors' calculations.

With respect to receiving at least two tetanus inoculations, there is no behavioral difference between uneducated women and those who have completed primary school. This implies that, when we take cluster-specific factors into account, women's education up to primary level has no impact on the tetanus inoculations they receive. Women who have completed up to middle or high school are more likely to receive tetanus toxoid injections compared to uneducated women. However, the magnitude of the effect is substantially lower than in the simple IV probit model.

In order to verify the results obtained from the cross-sectional and fixed effects models for overall Pakistan data, we conduct a series of robustness checks. A linear probability model for the two-stage IV probit model with district fixed effects is used to confirm the findings of the cluster fixed effects model. The results of both models are consistent with the primary models and are available on request.

The fixed effects model reveals some interesting findings on the impact of the husband's education on a woman's maternal health-seeking behavior. The effect of all levels of education – except secondary and higher education – becomes insignificant once community-specific characteristics are controlled for. While the effect of education completed up to secondary or higher level remains significant, there is a substantial decline in the size of this effect. An unexpected finding is that, once cluster-specific factors are controlled for, a woman whose spouse has completed only primary school is less likely to visit a healthcare facility at least three times seeking antenatal care compared to a woman whose spouse is uneducated.

The fixed effects model also indicates that, prior to controlling for cluster-specific characteristics, the coefficient of health knowledge is attenuated. Table 5 shows that, while the effect of women's health knowledge on their maternal health-seeking behavior remains significant, the magnitude of this influence has declined.

Once we control for heterogeneity within a sampling unit, there is no difference between the maternal health-seeking behavior of older and younger women, except in using safe delivery care. Similarly, women's maternal history does not influence their maternal healthcare use. The probability of women seeking safe delivery care is not affected by whether their pregnancy is planned or mistimed. This implies that the effect of planned pregnancy on women's use of safe delivery care is indeed confounded by area-specific characteristics.

Once these characteristics are controlled for, women are inclined to seek maternal healthcare services irrespective of whether their pregnancy is planned or unplanned, due to the risks associated with childbirth. The fixed effects model also shows that women who are married to the household head are less likely to receive safe delivery and postnatal care. Another notable finding is that consanguineous marriages have absolutely

no effect on women's maternal health-seeking behavior, once cluster-specific characteristics are controlled for.

Another key result of the fixed effects model is that the heterogeneity within sampling units seems to have the largest confounding effect on tetanus inoculations received by pregnant women in Pakistan. The results of cluster fixed effects show that women's age, primary schooling or employment status, previous fetal loss or stillbirth, and spouse's employment status do not influence their probability of receiving tetanus inoculations.

6. Conclusion and Policy Recommendations

This study has aimed to identify the socioeconomic determinants of maternal health-seeking behavior among women in Pakistan, using the IV approach. In the wake of binary dependent variables, we have used an IV probit model and analyzed a number of individual, household, and community-specific characteristics in relation to women.

Among individual characteristics, women's educational attainment as well as their spouses' education, women's health knowledge, and children's birth order emerge as key determinants of maternal health-seeking behavior. The results of both the cross-sectional as well as fixed effects models confirm that women's formal schooling significantly influences their maternal health-seeking behavior.

These effects of education are not, however, uniform across different levels of women's education, the net effect being highest for women who have completed secondary or higher education. Interestingly, we find that women with any level of formal schooling (even primary school) are more likely to seek maternal healthcare, thus reiterating the importance of maternal education.

On the other hand, whether a woman's husband has completed up to middle or high school emerges as a significant factor in maternal healthcare use. This implies that primary schooling alone is not enough to transform a spouse's attitude toward modern medicine. Government policies should, therefore, aim to give men more incentive to complete beyond primary school, achieve a better standard of living, and develop a more positive perception of modern medicine.

Another important finding is that women's health knowledge strongly influences their maternal health-seeking behavior, even after controlling for education. This implies that, while formal education might equip women with the tools to obtain health knowledge, there are also other sources that provide information on health, such as radio, television, and other mass media.

The results confirm that health knowledge can be acquired at any stage in life. Information on maternal healthcare broadcasted to the public – particularly women of childbearing age – through mass media public service programs could help increase women's awareness of maternal healthcare as well as their desire to learn how to better handle maternity. This should, in turn, encourage them to actively seek assistance from healthcare centers.

Our data shows that working women tend to come from less affluent households, where the decision to work is more likely to be rooted in need. These findings confirm that, in a society characterized by complex family structures, women's empowerment is reflected in their involvement in decision making. The results for spousal discussion of the place of delivery support this argument.

Education increases women's knowledge, capabilities, and self-confidence, all of which are also empowering. In turn, women's increased self-awareness modifies the traditional balance of power in family structures and helps women assume greater responsibility for their health-seeking behavior. While women's empowerment is an important determinant of their maternal health-seeking behavior, it is beyond the scope of this study and could prove an interesting area of future research.

As expected, the household's socioeconomic status, represented by the wealth index, emerges as one of the most important determinants of maternal healthcare utilization by women. This is not a novel finding, but it does underscore the strong effect of poverty on women's maternal health-seeking behavior. Household socioeconomic status does not appear to determine the tetanus inoculations received by women. The latter variable seems to be independent of most individual and household characteristics. Studying which factors determine the uptake of inoculations could, therefore, be an area of further research.

The Lady Health Workers Program is one of Pakistan's most extensive healthcare programs. Most women might receive tetanus shots through the door-to-door service that is part of lady health workers' stipulated job description. The results point to the importance of supply-side factors. Provided that lady health workers are the most pervasive source of inoculations for rural women, improving their outreach could help improve women's health significantly.

Several important policy implications emerge from these empirical findings. First, having established that women's health knowledge is influenced by, but not necessarily restricted to, their formal schooling, transmitting information on safe maternal healthcare practices through mass media (such as radio or television) may play an important role in this regard. Moreover, public programs could help effectively increase health knowledge among women of childbearing age and improve their maternal health-seeking behavior.

Second, the negative effect of women's employment and the fact that most working women are uneducated may reflect lower earnings. Skills-oriented training programs targeting less educated or illiterate women could increase their productivity and help generate better employment opportunities for them.

Finally, the differences in women's maternal health-seeking behavior across Punjab, Sindh, and KP imply that there is a need to improve the accessibility and quality of services available in these areas. In order to formulate effective policies at the provincial level, the social and cultural context of each province should be taken into account. The passage of the 18th Amendment has made the provinces responsible for designing and implementing health programs in line with their own needs. Nonetheless, the supervisory role of the federal government remains crucial.

References

- Afzal, U. (2013). What matters in child health: An instrumental variable analysis. *Child Indicators Research*, 6(4), 673–693.
- Agha, S., & Carton, T. W. (2011). Determinants of institutional delivery in rural Jhang, Pakistan. *International Journal for Equity in Health*, 10(31), 1–12.
- Ahmed, S., & Mosley, W. H. (2002). Simultaneity in the use of maternal-child healthcare and contraceptives: Evidence from developing countries. *Demography*, 39(1), 75–93.
- Ali, M., Bhatti, M. A., & Kuroiwa, C. (2008). Challenges in access to and utilization of reproductive healthcare in Pakistan. *Journal of Ayub Medical College Abbottabad*, 20(4), 3–7.
- Amin, R., Shah, N. M., & Becker, S. (2010). Socioeconomic factors differentiating maternal and child health-seeking behavior in rural Bangladesh: A cross-sectional analysis. *International Journal for Equity in Health*, 9(9), 1–11.
- Anderson, R. M. (1968). *Behavioral model of families' use of health services* (Research Series No. 25). Chicago, IL: University of Chicago, Center of Health Administration Studies.
- Andersen, R. M. (1995). Revisiting the behavioral model and access to medical care: Does it matter? *Journal of Health and Social Behavior*, 36(1), 1–10.
- Anderson, J. G., & Bartkus, D. E. (1973). Choice of medical care: A behavioral model of health and illness behavior. *Journal of Health and Social Behavior*, 14(4), 348–362.
- Anwar, M., Green, J., & Norris, P. (2012). Health-seeking behavior in Pakistan: A narrative review of the existing literature. *Public Health*, 126(6), 507–517.
- Aslam, M., & Kingdon, G. G. (2012). Parental education and child health: Understanding the pathways of impact in Pakistan. *World Development*, 40(10), 2014–2032.
- Becker, S., Fonseca-Becker, F., & Schenck-Yglesias, C. (2006). Husbands' and wives' reports of women's decision-making power in western Guatemala and their effects on preventive health behaviors. *Social Science and Medicine*, 62(9), 2313–2326.

- Bhatia, J. C., & Cleland, J. (1995). Determinants of maternal care in a region of South India. *Health Transition Review*, 5, 127–142.
- Bloom, S. S., Wypij, D., & Das Gupta, M. (2001). Dimensions of women's autonomy and the influence on maternal healthcare utilization in a north Indian city. *Demography*, 38(1), 67–78.
- Celik, Y., & Hotchkiss, D. R. (2000). The socioeconomic determinants of maternal healthcare utilization in Turkey. *Social Science and Medicine*, 50(12), 1797–1806.
- Chauhan, A. (2012). Antenatal care among currently married women in Rajasthan, India. *Asian Pacific Journal of Tropical Disease*, 2(2), 617–623.
- Das Gupta, M., Mansuri, G., Sinha, N., & Vishwanath, T. (2007, March). *Overcoming gender-based constraints to utilization of maternal and child health services in Pakistan: The role of the doorstep delivery system*. Paper presented at the Annual Meeting of the Population Association of America, New York, NY.
- Elo, I. T. (1992). Utilization of maternal healthcare services in Peru: The role of women's education. *Health Transition Review*, 2(1), 49–69.
- Furuta, M., & Salway, S. (2006). Women's position within the household as a determinant of maternal healthcare use in Nepal. *International Family Planning Perspectives*, 32(1), 17–27.
- Gage, A. J., & Calixte, M. G. (2006). Effects of the physical accessibility of maternal health services on their use in rural Haiti. *Population Studies*, 60(3), 271–288.
- Glewwe, P. (1999). Why does mother's schooling raise child health in developing countries? Evidence from Morocco. *Journal of Human Resources*, 34(1), 124–159.
- Gyimah, S. O., Takyi, B. K., & Addai, I. (2006). Challenges to the reproductive-health needs of African women: On religion and maternal health utilization in Ghana. *Social Science and Medicine*, 62(12), 2930–2944.
- Hassan, H., Jokhio, A. H., Winter, H., & MacArthur, C. (2012). Safe delivery and newborn care practices in Sindh, Pakistan: A community-based investigation of mothers and health workers. *Midwifery*, 28(4), 466–471.

- Hou, X., & Ma, N. (2011). *Empowering women: The effect of women's decision-making power on reproductive health services uptake: Evidence from Pakistan* (Policy Research Working Paper No. 5543). Washington, DC: World Bank.
- Khan, A. (1999). Mobility of women and access to health family planning services in Pakistan. *Reproductive Health Matters*, 7(14), 39–48.
- Kovsted, J., Pörtner, C. C., & Tarp, F. (2002). Child health and mortality: Does health knowledge matter. *Journal of African Economies*, 11(4), 542–560.
- LeVine, R. A., LeVine, S. E., Rowe, M. L., & Schnell-Anzola, B. (2004). Maternal literacy and health behavior: A Nepalese case study. *Social Science and Medicine*, 58(4), 863–877.
- McAlister, C., & Baskett, T. F. (2006). Female education and maternal mortality: A worldwide survey. *Journal of Obstetrics and Gynaecology Canada*, 28(11), 983–990.
- McTavish, S., Moore, S., Harper, S., & Lynch, J. (2010). National female literacy, individual socioeconomic status, and maternal healthcare use in sub-Saharan Africa. *Social Science and Medicine*, 71(11), 1958–1963.
- Midhet, F., Becker, S., & Berendes, H. W. (1998). Contextual determinants of maternal mortality in rural Pakistan. *Social Science and Medicine*, 46(12), 1587–1598.
- Mullany, B. C., Becker, S., & Hindin, M. J. (2007). The impact of including husbands in antenatal health education services on maternal health practices in urban Nepal: Results from a randomized control trial. *Health Education Research*, 22(2), 166–176.
- Mumtaz, Z., & Salway, S. M. (2007). Gender, pregnancy and the uptake of antenatal care services in Pakistan. *Sociology of Health and Illness*, 29(1), 1–26.
- Nisar, N., & White, E. (2003). Factors affecting utilization of antenatal care among reproductive age group women (15–49 years) in an urban squatter settlement of Karachi. *Journal of Pakistan Medical Association*, 53(2), 47–53.
- Sepehri, A., Sarma, S., Simpson, W., & Moshiri, S. (2008). How important are individual, household and commune characteristics in

- explaining utilization of maternal health services in Vietnam? *Social Science and Medicine*, 67(6), 1009–1017.
- Shariff, A., & Singh, G. (2002). *Determinants of maternal healthcare utilization in India: Evidence from a recent household survey* (Working Paper No. 85). New Delhi: National Council of Applied Economic Research.
- Singh, P. K., Rai, R. K., Alagarajan, M., & Singh, L. (2012). Determinants of maternity care services utilization among married adolescents in rural India. *PLOS ONE*, 7(2), e31666.
- Thomas, D., Strauss, J., & Henriques, M.-H. (1991). How does mother's education affect child height? *Journal of Human Resources*, 26(2), 183–211.
- Tura, G., & Gebremariam, A. (2008). Safe delivery service utilization in Metekel zone, northwest Ethiopia. *Ethiopia Journal of Health Sciences*, 17(4), 213–222.
- Webb, P., & Block, S. (2003). *Nutrition knowledge and parental schooling as inputs to child nutrition in the long and short run* (Nutrition Working Paper No. 3). Washington, DC: Development Alternatives, Inc.
- Winkvist, A., & Akhtar, H. Z. (1997). Images of health and healthcare options among low-income women in Punjab, Pakistan. *Social Science and Medicine*, 45(10), 1483–1491.
- Woldemicael, G. (2007). *Do women with higher autonomy seek more maternal and child healthcare? Evidence from Ethiopia and Eritrea* (Working Paper No. 2007-035). Rostock: Max Planck Institute for Demographic Research.
- World Health Organization. (2006). *Provision of effective antenatal care: Integrated management of pregnancy and childbirth (IMPAC)* (Standards for Maternal and Neonatal Care No. 1.6). Available from http://www.who.int/reproductivehealth/publications/maternal_perinatal_health/effective_antenatal_care.pdf
- World Health Organization. (2013). *Data and statistics* [Webpage]. Retrieved from <http://www.who.int/research/en/>

Appendix

**Table A1: First-stage regression results of health knowledge
(endogenous covariate)**

Explanatory variable	Coefficient
Instrumental variable	
Mass media exposure	
Heard a family planning message on radio or TV in the last month	0.0923*** (0.0183)
Individual characteristics	
Age (years)	
< 25 (base case)	
25–34	0.0201 (0.0204)
35–49	-0.0279 (0.0288)
Education	
None or less than primary (base case)	
Primary but below middle	-0.0120 (0.0289)
Middle but below secondary	-0.00990 (0.0329)
Secondary and above	0.0327 (0.0289)
Working woman	0.133*** (0.0213)
Child's birth order	
1 (base case)	
2 to 3	0.0354* (0.0210)
4 to 6	0.0551** (0.0259)
7+	0.102*** (0.0341)
Previous fetal loss or stillbirth	0.111*** (0.0207)
Previous fetal loss or stillbirth * working woman	-0.0648* (0.0368)
Planned pregnancy	-0.0622*** (0.0183)
Husband's education	
None or less than primary (base case)	

Explanatory variable	Coefficient
Primary but below middle	0.00774 (0.0251)
Middle but below secondary	0.0244 (0.0238)
Secondary and above	0.0108 (0.0214)
Husband has skilled employment	-0.0374** (0.0165)
Discussed place of delivery with spouse	0.0866*** (0.0164)
Husband is a blood relation	0.0287* (0.0156)
Wife of household head	0.0199 (0.0179)
Household characteristics	
Household wealth (index)	-0.0193* (0.0108)
Number of household members	0.000750 (0.00164)
Community characteristics	
Place of residence	
Region	
Urban (reference)	
Rural	-0.0061 (0.0250)
Punjab (reference)	
Sindh	0.270*** (0.0254)
KP	-0.0159 (0.0323)
Constant	0.251*** (0.0427)
Number of observations	4,174
F (25, 949)	18.43
R-squared	0.1323

Notes: * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$.

Robust standard errors at cluster level are given in parentheses below estimates.

Source: Authors' calculations based on PDHS 2006/07 data.

Table A2: Proportion of working women, by socioeconomic status

Wealth	Women working		Women not working	
	Number	Percentage	Number	Percentage
Low	338	38	502	17
Medium low	250	25	587	20
Medium	247	17	593	20
Medium high	181	12	659	21
High	127	7	709	21

Source: Authors' calculations based on PDHS 2006/07 data.

Table A3: Working women, by employment type

Employment type	Number	Percentage
Skilled work	609	13.62
Unskilled work	3,862	86.38
Total	4,471	100.00

Source: Authors' calculations based on PDHS 2006/07 data.

Lahore School of Economics
Centre for Research in Economics & Business

Recent Working Papers

No. 03-15

The Impact of Remittances and Parental Absence on Children's Wellbeing in Rural Punjab
Nida Jamil

No. 02-15

Natural Disasters and Social Capital: Evidence from a Field Experiment in Pakistan
Uzma Afzal, Ginger Turner and Farah Said

No. 01-15

Caste, Social Exclusion, and Opportunities for Education in Rural Punjab
Tayyaba Tamim, Marriam Musa, Hana Tariq and Shahzeb Rathore

No. 04-14

Education, Employment, and Women's Say in Household Decision-Making in Pakistan
Duryab Fatima

No. 03-14

The Impact of Parents' Subjective Aspirations on Schooling Investment in Rural Punjab
Anam Ashraf

No. 02-14

Attitudes Towards Risk in the Wake of a Rare Event: Evidence from Pakistan
Farah Said, Uzma Afzal and Ginger Turner

No. 01-14

Resource Misallocation and Aggregate Productivity in Punjab
Muhammad Haseeb and Theresa Thompson Chaudhry

No. 04-13

Labor Pooling as a Determinant of Industrial Agglomeration
Najam uz Zehra Gardezi

No. 03-13

The Effects of Agglomeration on the Formation and Scale of Operation of New Firms
Maryiam Haroon

No. 02-13

Agglomeration and Firm Turnover
Marjan Nasir

No. 01-13

Determinants of School Choice: Evidence from Rural Punjab, Pakistan
Hamna Ahmed, Sahar Amjad, Masooma Habib and Syed Ahsan Shah

No. 03-12

The Effects of External Migration on Enrolments, Accumulated Schooling, and Dropouts in Punjab
Rabia Arif and Azam Chaudhry

No. 02-12

The Determinants of Child Health and Nutritional Status in Punjab: An Economic Analysis
Uzma Afzal

No. 01-12

Investigating the Proposed Changes to Pakistan's Corporate Bankruptcy Code
Ali Hasanain and Syed Ahsan Ahmad Shah

Policy Papers

No. 01-11

Pakistan and Lessons from East Asia: Growth, Equity, and Governance
Khalid Ikram

No. 01-10

A Strategy for Reversing Pakistan's Dismal Export Performance
Hamna Ahmed, Mahreen Mahmud, Naved Hamid and Talal-Ur-Rahim

These papers can be accessed at: www.creb.org.pk

The Lahore School of Economics (established in 1993) is one of Pakistan's leading centres of learning for teaching and research in economics, finance and business administration. Its objectives are (i) to train young Pakistanis as professional economists, finance managers, accountants, financial analysts, bankers, and business executives, and (ii) to undertake research in economics, management, finance, and banking to deepen the understanding of major facts, issues, and policies.

The Centre for Research in Economics and Business (CREB) is an independent research centre at the Lahore School of Economics. CREB's mission is to conduct and facilitate research, coordinate and manage the Lahore School's postgraduate program, and promote discussion on policy issues facing Pakistan. The research focus at CREB is on the management of the Pakistan economy, income distribution and poverty, and the role of the modern services sector in the area of economics; and financial markets in the area of business management.

The Lahore School's publication program comprises the Lahore Journal of Economics, Lahore Journal of Policy Studies, Lahore Journal of Business, a Text Book Series, Lahore School Case Study Journal, the CREB Working Paper Series, and CREB Policy Paper Series. The program encourages both in-house and external contributors.



Lahore School of Economics

**Intersection Main Boulevard Phase VI, DHA and Burki Road
Burki Lahore 53200, Pakistan**